SLOPE AND ITS ROLE IN DESIGN FOR GENERALIZATION

Xinran Miao

University of Wisconsin-Madison

Joint work with Jiwei Zhao and Hyunseung Kang

ACIC 2025



"Science is about generalization, and generalization requires that conclusions obtained in the laboratory be transported and applied elsewhere" [Pearl and Bareinboim 2014]

"Science is about generalization, and generalization requires that conclusions obtained in the laboratory be transported and applied elsewhere" [Pearl and Bareinboim 2014]

Population	X	0	Goal
Q (Target)	√		$\psi(\mathbf{O})$
P (Source)	√	\checkmark	$\psi(\mathbf{Q}_{O,X})$

A common task: studying a summary of a target distribution $Q_{O,X}$, denoted as $\psi(Q_{O,X})$, where only the marginal Q_X can be identified.

"Science is about generalization, and generalization requires that conclusions obtained in the laboratory be transported and applied elsewhere" [Pearl and Bareinboim 2014]

Population	X	0	Goal		A common task: studying a summary of a target distribution					
Q (Target)	\		ab(O)	$Q_{0,X}$, denoted as $\psi(Q_{0,X})$, where only the marginal Q_X can be					
P (Source)	\	\checkmark	ψ(Ψ),X)	ident	ified.				
				X	$O \qquad P_{O,X} Q_X \psi(Q_{O,X})$					
Effect gener	raliza	tion		Base	seline Potential outcome Peru India $E_{Q_{O,X}}(O)$					
LLM [Mirza	deh (et al.	2024]	Mat	ath question Correct/Incorrect (1/0) Sophie Julia $pr_{Q_O}(O = 1)$					



Q = India

"Science is about generalization, and generalization requires that conclusions obtained in the laboratory be transported and applied elsewhere" [Pearl and Bareinboim 2014]

Population	X	0	Goal		A co	ommon task: studying a su	immary o	f a targe	et distribution	
Q (Target)	√		abl O)	$Q_{O,X}$, denoted as $\psi(Q_{O,X})$, whe	ere only th	ne marg	inal Q_X can be	
P (Source)	\	✓	Ψ(Ψ),X)	identified.					
				X	$O \qquad P_{O,X} Q_X \psi(Q_{O,X})$					
Effect gener	raliza	tion		Baseline	seline Potential outcome Peru India $E_{Q_{0,X}}(O)$					
LLM [Mirza	deh	et al.	2024]	Math qu	ath question Correct/Incorrect (1/0) Sophie Julia $\operatorname{pr}_{Q_O}(O=1)$					

P =Sophie

Sophie gets toys for her nephew... **31** blocks...**8** stuffed animals...**Sophie** recently bought bouncy balls... total number of toys is **62**. **How many** bouncy balls?

Q =Julia

Julia gets toys for her nephew... 31 blocks...8 stuffed animals...Julia recently bought bouncy balls... total number of toys is 62. How many bouncy balls?

"Science is about generalization, and generalization requires that conclusions obtained in the laboratory be transported and applied elsewhere" [Pearl and Bareinboim 2014]

Population	X	0	Goal		A co	ommon task: studying a su	immary o	f a targe	et distribution	
Q (Target)	√		ab(0))	$Q_{O,X}$, denoted as $\psi(Q_{O,X})$, where only the marginal Q_X can be					
P (Source)	\	\checkmark	$\psi(\mathbf{u})$),X)	identified.					
-										
				X	O $P_{O,X}$ Q_X $\psi(Q_{O,X})$				$\psi(Q_{O,X})$	
Effect gener	raliza	tion		Baseline	seline Potential outcome Peru India $E_{Q_{O,X}}(O)$					
LLM [Mirza	adeh	et al.	2024]	Math qu	ath question Correct/Incorrect (1/0) Sophie Julia $pr_{Q_0}(O = 1)$					

To borrow from $P_{O,X}$, the conditional exchangeability assumption is widely adopted:

$$Q_{O|X}(\cdot \mid x) = P_{O|X}(\cdot \mid x), \text{ a.e. } Q_X.$$
(1)

For example,

$$\psi(Q_{O,X}) = \mathcal{E}_{Q_O}(O)$$

= $\mathcal{E}_{Q_X} \left[\mathcal{E}_{Q_O \mid X}(O \mid X) \right]$
= $\mathcal{E}_{Q_X} \left[\mathcal{E}_{P_O \mid X}(O \mid X) \right].$

"Science is about generalization, and generalization requires that conclusions obtained in the laboratory be transported and applied elsewhere" [Pearl and Bareinboim 2014]

Population	X	0	Goal		A co	mmon task: studying a su	immary o	f a targe	et distribution	
Q (Target)	 Image: A start of the start of		ab(0))	$Q_{O,X}$, denoted as $\psi({\it Q}_{O,X})$, whe	ere only tl	he marg	inal Q_X can be	
P (Source)	\	\checkmark	$\psi(\mathbf{u})$),X)	ident	ified.				
				X	O $P_{O,X}$ Q_X $\psi(Q_{O,X})$				$\psi(Q_{O,X})$	
Effect gener	raliza	tion		Baseline	seline Potential outcome Peru India $E_{Q_{O,X}}(O)$					
LLM [Mirza	adeh	et al.	2024]	Math ques	ath question Correct/Incorrect (1/0) Sophie Julia $pr_{Q_0}(O = 1)$					

To borrow from $P_{O,X}$, the conditional exchangeability assumption is widely adopted:

$$Q_{O|X}(\cdot \mid x) = P_{O|X}(\cdot \mid x), \text{ a.e. } Q_X.$$
⁽¹⁾

For example,

$$\psi(Q_{O,X}) = \mathcal{E}_{Q_O}(O)$$

= $\mathcal{E}_{Q_X} \left[\mathcal{E}_{Q_O|X}(O \mid X) \right]$
= $\mathcal{E}_{Q_X} \left[\mathcal{E}_{P_{O|X}}(O \mid X) \right].$

However, the conditional exchengeability is made on unobserved data and cannot be verified.

<u>Our work</u>: If conditional exchengeability is **implausible**, should we (a) still borrow from P_{O,X}? (b) still report the mean?
 Are all **source distributions** (P_{O,X}) and **estimands** (ψ) equally "good/bad" when conditional exchengeability doesn't hold?

• <u>Our work</u>: If conditional exchengeability is **implausible**, should we (a) still borrow from $P_{O,X}$? (b) still report the mean?

Are all **source distributions** ($P_{O,X}$) and **estimands** (ψ) equally "good/bad" when conditional exchengeability doesn't hold?

Let $Q_{O,X}^{\gamma}$ be the distribution of $Q_{O,X}$ when conditional exchangeability is "violated" by some magnitude $\gamma \in \mathbb{R}$, where $\gamma = 0$ implies conditional exchangeability.



• <u>Our work</u>: If conditional exchangeability is **implausible**, should we (a) still borrow from $P_{O,X}$? (b) still report the mean?

Are all **source distributions** ($P_{O,X}$) and **estimands** (ψ) equally "good/bad" when conditional exchengeability doesn't hold?

- Let $Q_{O,X}^{\gamma}$ be the distribution of $Q_{O,X}$ when conditional exchangeability is "violated" by some magnitude $\gamma \in \mathbb{R}$, where $\gamma = 0$ implies conditional exchangeability.
- ▶ We propose a scalar summary, SLOPE. Higher $|SLOPE| \Rightarrow$ more sensitive to violations.



• <u>Our work</u>: If conditional exchengeability is **implausible**, should we (a) still borrow from $P_{O,X}$? (b) still report the mean?

Are all **source distributions** ($P_{O,X}$) and **estimands** (ψ) equally "good/bad" when conditional exchengeability doesn't hold?

- Let $Q_{O,X}^{\gamma}$ be the distribution of $Q_{O,X}$ when conditional exchangeability is "violated" by some magnitude $\gamma \in \mathbb{R}$, where $\gamma = 0$ implies conditional exchangeability.
- We propose a scalar summary, SLOPE. Higher $|SLOPE| \Rightarrow$ more sensitive to violations.



• <u>Our work</u>: If conditional exchengeability is **implausible**, should we (a) still borrow from $P_{O,X}$? (b) still report the mean?

Are all **source distributions** ($P_{O,X}$) and **estimands** (ψ) equally "good/bad" when conditional exchengeability doesn't hold?

- Let $Q_{O,X}^{\gamma}$ be the distribution of $Q_{O,X}$ when conditional exchangeability is "violated" by some magnitude $\gamma \in \mathbb{R}$, where $\gamma = 0$ implies conditional exchangeability.
- ▶ We propose a scalar summary, SLOPE. Higher $|SLOPE| \Rightarrow$ more sensitive to violations.



OUTLINE

1 Proposal: SLOPE

2 Data Application

3 Summary

NOTATION AND REVIEW OF KEY CONCEPTS

- ▶ Target distribution: $Q_{O,X}$ of (O, X) where $O \in \mathbb{R}$ and $X \in \mathcal{X} \subset \mathbb{R}^d$.
- Scientific interest: a low dimensional $\psi(Q_{O,X})$ with $\psi(\cdot)$ defined by investigator.
- We only have access to Q_X and a related source distribution $P_{O,X}$.

In order to identify the functional $\psi(Q_{O,X})$ using the random sample from $P_{O,X}$ and Q_X , two assumptions are often imposed.

Assumption 1 (Overlap)

 Q_X is absolute continuous with respect to P_X .

Assumption 2 (Conditional Exchangeability)

 $Q_{O|X}(\cdot \mid x) = P_{O|X}(\cdot \mid x)$ almost everywhere Q_X .

Under Assumptions 1-2, the target functional can be identified:

$$\psi(Q_{O,X}) = \psi(Q_X \times Q_{O|X}) = \psi(Q_X \times P_{O|X}).$$

Example functionals include the mean ψ^{MEAN} , median ψ^{MED} , and OLS coefficient ψ^{OLS} .

Conditional exchengeability is made on the unobserved data and cannot be empirically verified [Nguyen et al. 2017; Nie, Imbens, and Wager 2021; Huang 2024]. Assuming it blindly can lead to distorted conclusions [Jin, Egami, and Rothenhäusler 2024].

Conditional exchengeability is made on the unobserved data and cannot be empirically verified [Nguyen et al. 2017; Nie, Imbens, and Wager 2021; Huang 2024]. Assuming it blindly can lead to distorted conclusions [Jin, Egami, and Rothenhäusler 2024].

We follow the Robins' sensitivity model [Robins, Rotnitzky, and Scharfstein 2000; Franks, D'Amour, and Feller 2020; Nabi et al. 2024] and assume that the distribution $Q_{O|X}$ is shifted from $P_{O|X}$ by an exponential tilting term $\exp(\gamma o)$:

$$\frac{\mathrm{d}Q_{O|X}^{\gamma}(o \mid x)}{\mathrm{d}P_{O|X}(o \mid x)} \propto \exp(\gamma \cdot o), \tag{2}$$

where we assume $\int \exp(\gamma o) dP_{O|X}(o, x) < \infty$.

Conditional exchengeability is made on the unobserved data and cannot be empirically verified [Nguyen et al. 2017; Nie, Imbens, and Wager 2021; Huang 2024]. Assuming it blindly can lead to distorted conclusions [Jin, Egami, and Rothenhäusler 2024].

We follow the Robins' sensitivity model [Robins, Rotnitzky, and Scharfstein 2000; Franks, D'Amour, and Feller 2020; Nabi et al. 2024] and assume that the distribution $Q_{O|X}$ is shifted from $P_{O|X}$ by an exponential tilting term $\exp(\gamma o)$:

$$\frac{\mathrm{d}Q_{O|X}^{\gamma}(o \mid x)}{\mathrm{d}P_{O|X}(o \mid x)} \propto \exp(\gamma \cdot o), \tag{2}$$

where we assume $\int \exp(\gamma o) dP_{O|X}(o, x) < \infty$.

- $\gamma \in \mathbb{R}$ is a sensitivity parameter that measures the unobserved differences between $Q_{O|X}$ and $P_{O|X}$.
- ▶ $\gamma = 0 \implies$ conditional exchangeability holds.
- $\gamma \neq 0 \Longrightarrow$ conditional exchangeability is violated.
- $\blacktriangleright \ \gamma$ cannot be identified from data.

Conditional exchengeability is made on the unobserved data and cannot be empirically verified [Nguyen et al. 2017; Nie, Imbens, and Wager 2021; Huang 2024]. Assuming it blindly can lead to distorted conclusions [Jin, Egami, and Rothenhäusler 2024].

We follow the Robins' sensitivity model [Robins, Rotnitzky, and Scharfstein 2000; Franks, D'Amour, and Feller 2020; Nabi et al. 2024] and assume that the distribution $Q_{O|X}$ is shifted from $P_{O|X}$ by an exponential tilting term $\exp(\gamma o)$:

$$\frac{\mathrm{d}Q_{O|X}^{\gamma}(o \mid x)}{\mathrm{d}P_{O|X}(o \mid x)} \propto \exp(\gamma \cdot o), \tag{2}$$

where we assume $\int \exp(\gamma o) dP_{O|X}(o, x) < \infty$.

- $\gamma \in \mathbb{R}$ is a sensitivity parameter that measures the unobserved differences between $Q_{O|X}$ and $P_{O|X}$.
- $\gamma = 0 \Longrightarrow$ conditional exchangeability holds.
- ▶ $\gamma \neq 0 \implies$ conditional exchangeability is violated.
- γ cannot be identified from data.

Examples:

► If
$$P_{O|X} \sim N(\mu(X), \sigma^2(X))$$
, then (2) elicits a location shift, $Q_{O|X}^{\gamma} \sim N(\mu(X) + \gamma \sigma^2(X), \sigma^2(X))$.

• If *O* is binary, then γ has an odds ratio interpretation, $\exp(\gamma) = \frac{Q^{\gamma}(O = 1 \mid X) / \{1 - Q^{\gamma}(O = 1 \mid X)\}}{P(O = 1 \mid X) / \{1 - P(O = 1 \mid X)\}}$.

SLOPE

Let
$$Q_{O,X}^{\gamma} = Q_X \times Q_{O|X}^{\gamma}$$
.

Definition 1

The **S**ensitivity to **LO**cal **P**erturbation from **E**xchangeability (SLOPE) of a functional ψ with respect to the perturbation (2) is

$$\mathsf{SLOPE}(Q_{O,X}^0,\psi) = \lim_{\gamma \to 0} \frac{\psi(Q_{O,X}^\gamma) - \psi(Q_{O,X}^0)}{\gamma}, \tag{3}$$

provided the limit exists.

- SLOPE is a (usually) scalar summary to summarize sensitivity of an estimand when conditional exchangeability is violated in a particular "direction".
- ► Higher |SLOPE| ⇒ estimand changes drastically under local violations ⇒ more sensitive.



- ► SLOPE($Q_{O,X}^0, \psi$) depends on the distribution $Q_{O,X}^0 = Q_X \times P_{O|X}$ and the estimand ψ .
 - SLOPE can be applied to choose source $(P_{O|X})$ and estimand (ψ) in order to be less sensitive to violations of conditional exchangeability.



- SLOPE $(Q_{O,X}^0, \psi)$ depends on the distribution $Q_{O,X}^0 = Q_X \times P_{O|X}$ and the estimand ψ .
 - SLOPE can be applied to choose source $(P_{O|X})$ and estimand (ψ) in order to be less sensitive to violations of conditional exchangeability.



- SLOPE($Q_{O,X}^0, \psi$) depends on the distribution $Q_{O,X}^0 = Q_X \times P_{O|X}$ and the estimand ψ .
 - SLOPE can be applied to choose source $(P_{O|X})$ and estimand (ψ) in order to be less sensitive to violations of conditional exchangeability.



- SLOPE $(Q_{O,X}^0, \psi)$ depends on the distribution $Q_{O,X}^0 = Q_X \times P_{O|X}$ and the estimand ψ .
 - SLOPE can be applied to choose source $(P_{O|X})$ and estimand (ψ) in order to be less sensitive to violations of conditional exchangeability.



- SLOPE $(Q_{O,X}^0, \psi)$ depends on the distribution $Q_{O,X}^0 = Q_X \times P_{O|X}$ and the estimand ψ .
 - SLOPE can be applied to choose source $(P_{O|X})$ and estimand (ψ) in order to be less sensitive to violations of conditional exchangeability.
 - SLOPE is a population level quantity which reflects the design instead of estimation; it does not depend on any estimation procedure.
 - We think this way of summarizing sensitivity can be broadly helpful.



- SLOPE($Q_{O,X}^0, \psi$) depends on the distribution $Q_{O,X}^0 = Q_X \times P_{O|X}$ and the estimand ψ .
 - SLOPE can be applied to choose source $(P_{O|X})$ and estimand (ψ) in order to be less sensitive to violations of conditional exchangeability.
 - SLOPE is a population level quantity which reflects the design instead of estimation; it does not depend on any estimation procedure.
 - We think this way of summarizing sensitivity can be broadly helpful.
- Additional remarks on interpretation.
 - The unit of the SLOPE matches the unit of the estimand.
 - SLOPE is a first-order violation of conditional exchengeability, i.e.,

$$\psi(Q_{O,X}^{\gamma}) \approx \psi(Q_{O,X}^{0}) + \gamma \cdot \text{SLOPE}(Q_{O,X}^{0}, \psi).$$

SLOPE cannot identify the bias under violation (which is unknown).



Technical remarks:

- Finding the form of the SLOPE is not difficult; it's an exercise in calculus if we can freely interchange integration and differentiation.
- SLOPE can be consistently estimated under regularity conditions; not covered in this talk.
- SLOPE does not exist for every estimand; e.g., $sign_{Q_0}(O)$.



EXAMPLE: SLOPE FOR MEAN AND MEDIAN

Let $\sigma^2(X) = \operatorname{var}_{P_{O|X}}(O \mid X)$ be the conditional variance on the source population.

The SLOPE for the mean is

 $SLOPE(Q_{O,X}^0, \psi^{MEAN}) = E_{Q_X} \{ \sigma^2(X) \}.$ (SLOPE for Mean)

Example: SLOPE for Mean and Median

Let $\sigma^2(X) = \operatorname{var}_{P_{O|X}}(O \mid X)$ be the conditional variance on the source population.

The SLOPE for the mean is

 $SLOPE(Q_{O,X}^{0}, \psi^{MEAN}) = E_{Q_X} \{ \sigma^2(X) \}.$ (SLOPE for Mean)

Under the conditional Gaussian model, $P_{O|X} \sim N(\mu(X), \sigma^2(X))$, the SLOPE for the median is

$$SLOPE(Q_{O,X}^{0}, \psi^{MED}) = E_{Q_X} \left[\sigma^2(X) \cdot \underbrace{\frac{f_{P_{O|X}}(m_{1/2})}{E_{Q_X} \left\{ f_{P_{O|X}}(m_{1/2}) \right\}}}_{Weight} \right], \qquad (SLOPE \text{ for Median})$$

where $m_{1/2} = F_{Q_0^0}^{-1}(1/2)$ is the median and $f_{P_{O|X}}$ is the density with respect to some measure.

Example: SLOPE for Mean and Median

Let $\sigma^2(X) = \operatorname{var}_{P_{O|X}}(O \mid X)$ be the conditional variance on the source population.

The SLOPE for the mean is

 $SLOPE(Q_{O,X}^{0}, \psi^{MEAN}) = E_{Q_{X}}\{\sigma^{2}(X)\}.$ (SLOPE for Mean)

Under the conditional Gaussian model, $P_{O|X} \sim N(\mu(X), \sigma^2(X))$, the SLOPE for the median is

$$SLOPE(Q_{O,X}^{0},\psi^{MED}) = E_{Q_X} \left[\sigma^2(X) \cdot \frac{f_{P_{O|X}}(m_{1/2})}{\underbrace{E_{Q_X}\left\{f_{P_{O|X}}(m_{1/2})\right\}}} \right], \qquad (SLOPE \text{ for Median})$$
Weight

where $m_{1/2} = F_{Q_O^0}^{-1}(1/2)$ is the median and $f_{P_{O|X}}$ is the density with respect to some measure. SLOPE for location parameters depends heavily on heterogeneity of *O* in the source population. This matches with existing insights that homogeneous ATE is easier to generalize [Tipton 2014].

CONNECTION TO LITERATURE

Prior works are mostly on estimators rather than estimands.

- <u>Robust statistics</u>: "...robustness signifies insensitivity to small deviations from the assumptions..." [Huber 1981]
 - SLOPE is analogous to the influence function (IF) [Hampel 1974] as a local derivative.

$$SLOPE(Q_{O,X}^{0},\psi) = \lim_{\gamma \to 0} \frac{\psi(Q_{O,X}^{\gamma}) - \psi(Q_{O,X}^{0})}{\gamma}$$
$$IF(Q_{O,X}^{0},\psi,o,x) = \lim_{\varepsilon \to 0^{+}} \frac{\psi((1-\varepsilon)Q_{O,X}^{0} + \varepsilon \delta_{o,x}) - \psi(Q_{O,X}^{0})}{\varepsilon}$$

• For parameters defined through Z-estimation, SLOPE can be re-expressed with the IF (next slide).

CONNECTING SLOPE WITH IF IN Z-ESTIMATION

Suppose the finite-dimensional parameter ψ is defined through

$$E_{Q_{O,X}^{0}}\{s(O,X,\psi)\} = 0,$$
(4)

where *s* is a known function.

Theorem 1 (SLOPE and Influence Function for Z-Estimation)

The SLOPE for ψ *in* (4) *is*

$$\mathsf{SLOPE}(Q_{O,X}^0,\psi) = \iint \underbrace{\mathsf{IF}(O,X,\psi(Q_{O,X}^0))}_{\mathsf{Influence function}} \underbrace{\{O-\mu(X)\}}_{\mathsf{Direction of sensitivity}} dP_{O|X} dQ_X;$$

where $\mu(X) = E_{P_{O|X}}(O \mid X)$, and $IF(O, X, \psi(Q_{O,X}^0))$ is the influence function for $\psi(Q_{O,X}^0)$ under $Q_{O,X}^0$.

SLOPE is a recentered integral of the IF, where the re-centering is determined by $O - \mu(X)$.

OUTLINE

1 Proposal: SLOPE

2 Data Application

3 Summary

RUNNING EXAMPLE: TRANSPORTING POTENTIAL OUTCOMES

- [Banerjee et al. 2015] conducted a multi-site RCT to evaluate the effect of the "Graduation Program" on improving livelihoods of poor households in six countries (Ethiopia, Ghana, Honduras, India, Pakistan, Peru).
 - "...provides a holistic set of services, including the grant of a productive asset, to the poorest households in a village"
- ▶ We treat one country as the source (*P*) and another as the target (*Q*) and investigate the sensitivity to violations of conditional exchangeability between $Q_{O|X}$ and $P_{O|X}$.
- ► *X* is a categorical baseline measurement prior to intervention.
- O = Y(1) is the potential outcome under the intervention from "Graduation Program".
- ▶ $P = P_{O,X} = P_{Y(1),X}$: households in the source country (e.g., Honduras).
- $Q = Q_{O,X} = Q_{Y(1),X}$: households in the target country (e.g., Ethiopia), where we only observe the baseline Q_X .
- ▶ Identification of the SLOPE requires additional assumptions on *P*_{*O*,*X*} (SUTVA, strong ignorability).

▶ Outcome: log-transformed total asset index (standardized total asset value in terms of goats in each country). It ranges from −1.3 to 2.97.

Question: Which source country is the least sensitive?

Estimand (ψ)	Source $(P_{O X})$	Target (<i>Q_X</i>) Ethiopia
	Ethiopia	
	Ghana	0.13
Mean	Honduras	0.09
Mean	India	0.16
	Pakistan	0.16
	Peru	0.20

▶ Outcome: log-transformed total asset index (standardized total asset value in terms of goats in each country). It ranges from −1.3 to 2.97.

Question: Which source country is the least sensitive?

Estimond (a/a)	Source (D)	Target (<i>Q</i> _X)							
Estimanu (ψ)	Source $(P_{O X})$	Ethiopia	Ghana	Honduras	India	Pakistan	Peru		
	Ethiopia		0.15	0.18	0.15	0.14	0.13		
	Ghana	0.13		0.10	0.10	0.11	0.13		
Mean	Honduras	0.09	0.08		(0.07)	0.08	0.10		
Mean	India	0.16	0.15	0.16		0.16	0.16		
	Pakistan	0.16	0.15	0.14	0.15		0.16		
	Peru	0.20	0.21	0.21	0.21	0.21			

► Outcome: log-transformed total asset index (standardized total asset value in terms of goats in each country). It ranges from -1.3 to 2.97.

Question: Which source country is the least sensitive? **Answer**: Honduras.

Estimand (a/a)	Source (D, \cdot)			Target (Q_X)		
Estimatio (ψ)	Source $(r_{O X})$	Ethiopia	Ghana	Honduras	India	Pakistan	Peru
	Ethiopia		0.15	0.18	0.15	0.14	0.13
	Ghana	0.13		0.10	0.10	0.11	0.13
Mean	Honduras	0.09	0.08		0.07	0.08	0.10
mean	India	0.16	0.15	0.16		0.16	0.16
	Pakistan	0.16	0.15	0.14	0.15		0.16
	Peru	0.20	0.21	0.21	0.21	0.21	
	Ethiopia		0.15	0.18	0.15	0.14	0.13
	Ghana	0.11		0.10	0.09	0.09	0.11
Median	Honduras	0.07	0.06		0.05	0.06	0.07
	India	0.15	0.15	0.16		0.15	0.16
	Pakistan	0.16	0.15	0.14	0.15		0.16
	Peru	0.19	0.21	0.21	0.22	0.21	

► Outcome: log-transformed total asset index (standardized total asset value in terms of goats in each country). It ranges from -1.3 to 2.97.

Question: Which source country / estimand is the least sensitive? Answer: Honduras / Median.

Estimond (1/1)	Source (P, \cdot)			Target ($Q_{\chi})$		
Estimanu (ψ)	Source $(r_{O X})$	Ethiopia	Ghana	Honduras	India	Pakistan	Peru
	Ethiopia		0.15	0.18	0.15	0.14	0.13
	Ghana	0.13		0.10	0.10	0.11	0.13
Mean	Honduras	0.09	0.08		(0.07)	0.08	0.10
Mean	India	0.16	0.15	0.16		0.16	0.16
	Pakistan	0.16	0.15	0.14	0.15		0.16
	Peru	0.20	0.21	0.21	0.21	0.21	
	Ethiopia		0.15	0.18	0.15	0.14	0.13
	Ghana	0.11		0.10	0.09	0.09	0.11
Median	Honduras	0.07	0.06		0.05	0.06	0.07
, incuration	India	0.15	0.15	0.16		0.15	0.16
	Pakistan	0.16	0.15	0.14	0.15		0.16
	Peru	0.19	0.21	0.21	0.22	0.21	

The median is usually slightly better than the mean (i.e., has smaller SLOPE).

[Banerjee et al. 2015] reported the "physical health index", which is the average of z-scores of three variables:

- Did not miss work due to illness (work), Zwork between -4.3 and 7.0;
- Activities of daily living score (act), Z_{act} between -3.2 and 1.1;
- Perception of health status (perc), Z_{perc} between -2.3 and 4.5.

[Banerjee et al. 2015] reported the "physical health index", which is the average of z-scores of three variables:

- Did not miss work due to illness (work), Zwork between -4.3 and 7.0;
- Activities of daily living score (act), Z_{act} between -3.2 and 1.1;
- Perception of health status (perc), Z_{perc} between -2.3 and 4.5.

Question: Is there another way to define a physical health index so that it's less sensitive during generalization?

[Banerjee et al. 2015] reported the "physical health index", which is the average of z-scores of three variables:

- Did not miss work due to illness (work), Zwork between -4.3 and 7.0;
- Activities of daily living score (act), Z_{act} between -3.2 and 1.1;
- Perception of health status (perc), Z_{perc} between -2.3 and 4.5.

Question: Is there another way to define a physical health index so that it's less sensitive during generalization?

Consider linear combinations:

$$w_{\text{work}} \cdot Z_{\text{work}} + w_{\text{act}} \cdot Z_{\text{act}} + w_{\text{perc}} \cdot Z_{\text{perc}},$$

where the original index has $w_{work} = w_{act} = w_{perc} = 1/3$.

• On the simplex of $w = [w_{work}, w_{act}, w_{perc}]^T$, we want to find the weight that yields the smallest SLOPE.

<u>India (P)</u> \Rightarrow Ethiopia (Q)

 \star [$w_{work}, w_{act}, w_{perc}$]^T = [0.10, 0.55, 0.35]^T



 $\underline{\operatorname{Peru}\left(P\right)} \Rightarrow \operatorname{Ethiopia}\left(Q\right)$

$$\bigstar [w_{\text{work}}, w_{\text{act}}, w_{\text{perc}}]^{\text{T}} = [1, 0, 0]^{\text{T}}$$



OUTLINE

1 Proposal: SLOPE

2 Data Application

3 Summary

SUMMARY AND CONTACT

We propose a measure, sensitivity to local perturbation from exchangeability (SLOPE), that describes the change in a functional ψ with respect to a local perturbation (2):

$$\mathsf{SLOPE}(\boldsymbol{Q}_{O,X}^0,\psi) = \lim_{\gamma o 0} rac{\psi(\boldsymbol{Q}_{O,X}^\gamma) - \psi(\boldsymbol{Q}_{O,X}^0)}{\gamma},$$

provided the limit exists.

- SLOPE is a scalar summary for sensitivity to violation of the conditional exchengeability assumption.
- SLOPE guides two choices in design:
 - Choosing the source population *P*.
 - Choosing the estimand $\psi(\cdot)$.
- SLOPE connects to influence function (both conceptually and mathematically).
- Main limitations:
 - SLOPE assesses the local sensitivity (in a linear approximation).
 - SLOPE is not unit-less.

References I

- Andrews, Isaiah, Matthew Gentzkow, and Jesse M Shapiro (2017). "Measuring the sensitivity of parameter estimates to estimation moments". In: *The Quarterly Journal of Economics* 132.4, pp. 1553–1592.
- Banerjee, Abhijit et al. (2015). "A multifaceted program causes lasting progress for the very poor: Evidence from six countries". In: *Science* 348.6236, p. 1260799.
- Birmingham, Jolene, Andrea Rotnitzky, and Garrett M Fitzmaurice (2003). "Pattern-mixture and selection models for analysing longitudinal data with monotone missing patterns". In: Journal of the Royal Statistical Society Series B: Statistical Methodology 65.1, pp. 275–297.
- Cinelli, Carlos and Chad Hazlett (2020). "Making sense of sensitivity: Extending omitted variable bias". In: Journal of the Royal Statistical Society Series B: Statistical Methodology 82.1, pp. 39–67.
- Colnet, Bénédicte et al. (2023). "Risk ratio, odds ratio, risk difference... Which causal measure is easier to generalize?" In: *arXiv preprint arXiv:2303.16008*.
- Dahabreh, Issa J et al. (2022). "Global sensitivity analysis for studies extending inferences from a randomized trial to a target population". In: *arXiv preprint arXiv:2207.09982*.
- Ding, Peng and Tyler J VanderWeele (2016). "Sensitivity analysis without assumptions". In: *Epidemiology* 27.3, pp. 368–377.
- Franks, Alexander M, Alexander D'Amour, and Avi Feller (2020). "Flexible sensitivity analysis for observational studies without observable implications". In: Journal of the American Statistical Association 115.532, pp. 1730–1746.

References II

- Hampel, Frank R (1974). "The influence curve and its role in robust estimation". In: Journal of the american statistical association 69.346, pp. 383–393.
- Huang, Melody Y (2024). "Sensitivity analysis for the generalization of experimental results". In: *Journal of the Royal Statistical Society Series A: Statistics in Society* 187.4, pp. 900–918.
- Huber, P.J. (1981). *Robust statistics*. Wiley New York.
- Jin, Ying, Naoki Egami, and Dominik Rothenhäusler (2024). "Beyond Reweighting: On the Predictive Role of Covariate Shift in Effect Generalization". In: *arXiv preprint arXiv:2412.08869*.
- Mirzadeh, Iman et al. (2024). "Gsm-symbolic: Understanding the limitations of mathematical reasoning in large language models". In: *arXiv preprint arXiv:2410.05229*.
- Nabi, Razieh et al. (2024). "Semiparametric sensitivity analysis: unmeasured confounding in observational studies". In: *Biometrics* 80.4, ujae106.
- Nguyen, Trang Quynh et al. (2017). "Sensitivity analysis for an unobserved moderator in RCT-to-target-population generalization of treatment effects". In: The Annals of Applied Statistics, pp. 225–247.
- Nie, Xinkun, Guido Imbens, and Stefan Wager (2021). "Covariate balancing sensitivity analysis for extrapolating randomized trials across locations". In: *arXiv preprint arXiv:2112.04723*.
- **Oster, Emily (2019). "Unobservable selection and coefficient stability: Theory and evidence".** In: *Journal of Business & Economic Statistics* 37.2, pp. 187–204.

References III

- Pearl, Judea and Elias Bareinboim (2014). "External Validity: From Do-Calculus to Transportability Across Populations". In: *Statistical Science* 29.4, pp. 579–595.
- Robins, James M, Andrea Rotnitzky, and Daniel O Scharfstein (2000). "Sensitivity analysis for selection bias and unmeasured confounding in missing data and causal inference models". In: Statistical Models in Epidemiology, the Environment, and Clinical Trials. Springer, pp. 1–94.
- Rosenbaum, Paul R (2004). "Design sensitivity in observational studies". In: Biometrika 91.1, pp. 153–164.
- **Rotnitzky**, Andrea et al. (2001). "Methods for conducting sensitivity analysis of trials with potentially nonignorable competing causes of censoring". In: *Biometrics* 57.1, pp. 103–113.
- Tipton, Elizabeth (2014). "How generalizable is your experiment? An index for comparing experimental samples and populations". In: *Journal of Educational and Behavioral Statistics* 39.6, pp. 478–501.
- Tipton, Elizabeth and Robert B Olsen (2018). "A review of statistical methods for generalizing from evaluations of educational interventions". In: *Educational Researcher* 47.8, pp. 516–524.
- Troxel, Andrea B, Guoguang Ma, and Daniel F Heitjan (2004). "An index of local sensitivity to nonignorability". In: *Statistica Sinica*, pp. 1221–1237.
- Zhao, Qingyuan, Dylan S Small, and Bhaswar B Bhattacharya (2019). "Sensitivity analysis for inverse probability weighting estimators via the percentile bootstrap". In: Journal of the Royal Statistical Society Series B: Statistical Methodology 81.4, pp. 735–761.

Example Estimands $\psi(\cdot)$

• Outcome mean: $\psi^{MEAN}(Q_{O,X})$ which is

$$\psi^{\text{MEAN}}(Q_{O,X}) = \mathbb{E}_{Q_{O,X}}(O) = \mathbb{E}_{Q_X} \left[\mathbb{E}_{Q_{O|X}}(O) \right]$$
$$= \mathbb{E}_{Q_X} \left[\mathbb{E}_{P_{O|X}}(O) \right]$$
(By Assumptions 1-2).

• Outcome median: $\psi^{MED}(Q_{O,X})$ such that

$$1/2 = \int_{-\infty}^{\psi_{\text{MED}}} \mathrm{d}Q_O = \int \int_{-\infty}^{\psi_{\text{MED}}} \mathrm{d}Q_{O|X} \mathrm{d}Q_X$$
$$= \int \int_{-\infty}^{\psi_{\text{MED}}} \mathrm{d}P_{O|X} \mathrm{d}Q_X$$

(By Assumptions 1-2).

• OLS coefficient: $\psi^{OLS}(Q_{O,X})$ such that

$$0 = \mathbb{E}_{Q_{O,X}} \left[X X^{\mathrm{T}} \psi^{\mathrm{OLS}} - X O \right]$$

= $\mathbb{E}_{Q_X} \left[\mathbb{E}_{P_{O|X}} \left\{ X X^{\mathrm{T}} \psi^{\mathrm{OLS}} - X O \right\} \right]$ (By Assumptions 1-2).

EXAMPLES OF EXPONENTIAL TILT

$$\frac{\mathrm{d}Q_{O|X}^{\gamma}(o \mid x)}{\mathrm{d}P_{O|X}(o \mid x)} \propto \exp(\gamma \cdot o), \tag{2}$$

• If $P_{O|X} \sim N(\mu(X), \sigma^2(X))$, then (2) elicits a location shift

$$Q_{O|X}^{\gamma} \sim N\left(\mu(X) + \gamma\sigma^2(X), \sigma^2(X)\right).$$



$$\exp(\gamma) = \frac{Q^{\gamma}(O=1 \mid X) / \{1 - Q^{\gamma}(O=1 \mid X)\}}{P(O=1 \mid X) / \{1 - P(O=1 \mid X)\}}.$$





Why This Sensitivity Model?

$$\frac{\mathrm{d}Q_{O|X}^{\gamma}(o \mid x)}{\mathrm{d}P_{O|X}(o \mid x)} \propto \exp(\gamma \cdot o), \tag{2}$$

- Used in non-ignorable missing data [Robins, Rotnitzky, and Scharfstein 2000], unmeasured confounding in causal inference [Franks, D'Amour, and Feller 2020; Nabi et al. 2024], and effect generalization [Dahabreh et al. 2022].
- Simple to interprete (i.e., exponential tilting; odds ratio); can reparametrize γ to pseudo-R² [Franks, D'Amour, and Feller 2020].
- Does not place restrictions on observed data.
- Makes inference of the SLOPE (next slide) tractable (e.g., asymptotic normality).

CONNECTION TO LITERATURE

Prior works are mostly on estimators rather than estimands.

- <u>Robust statistics</u>: "...robustness signifies insensitivity to small deviations from the assumptions..." [Huber 1981]
 - SLOPE is analogous to the influence function (IF) [Hampel 1974] as a local derivative.

$$SLOPE(Q_{O,X}^{0},\psi) = \lim_{\gamma \to 0} \frac{\psi(Q_{O,X}^{\gamma}) - \psi(Q_{O,X}^{0})}{\gamma}$$
$$IF(Q_{O,X}^{0},\psi,o,x) = \lim_{\varepsilon \to 0^{+}} \frac{\psi((1-\varepsilon)Q_{O,X}^{0} + \varepsilon \delta_{o,x}) - \psi(Q_{O,X}^{0})}{\varepsilon}$$

- For parameters defined through Z-estimation, SLOPE can be re-expressed with the IF (next slide).
 <u>Sensitivity analysis</u>: our parametrization of Q^γ_{O|X} as deviations from Q⁰_{O,X} = P_{O|X} was proposed by [Robins, Rotnitzky, and Scharfstein 2000].
 - This line of work focuses on estimation/inference given $\gamma \neq 0$ [Rotnitzky et al. 2001; Birmingham, Rotnitzky, and Fitzmaurice 2003; Franks, D'Amour, and Feller 2020; Dahabreh et al. 2022; Nabi et al. 2024].
- Other related works: [Rosenbaum 2004; Troxel, Ma, and Heitjan 2004; Ding and VanderWeele 2016; Andrews, Gentzkow, and Shapiro 2017; Oster 2019; Cinelli and Hazlett 2020; Zhao, Small, and Bhattacharya 2019; Colnet et al. 2023; Jin, Egami, and Rothenhäusler 2024].

Example: SLOPE for Transporting the Mean

Suppose $O \in \mathbb{R}$ is an outcome variable. The target estimand is the outcome mean

$$\psi^{\scriptscriptstyle{\mathsf{MEAN}}}(Q_{O,X}) := \int O \mathrm{d} Q_O$$

Under the sensitivity model (2), the target functional with sensitivity parameter γ is

$$\psi^{\text{MEAN}}(Q_{O,X}^{\gamma}) = \int \frac{\int O \exp(\gamma O) dP_{O|X}}{\int \exp(\gamma O) dP_{O|X}} Q_X,$$
(5)

The SLOPE can be calculated via calculus:

$$\mathsf{SLOPE}(Q_{O,X}^0,\psi^{\mathsf{MEAN}}) = \frac{\partial \psi^{\mathsf{MEAN}}(Q_{O,X}^\gamma)}{\partial \gamma} \bigg|_{\gamma=0} = \mathsf{E}_{Q_X}\{\mathsf{var}_{P_{O|X}}(O \mid X)\} := \mathsf{E}_{Q_X}\{\sigma^2(X)\}.$$

EXAMPLE: SLOPE FOR TRANSPORTING THE MEAN

Suppose $O \in \mathbb{R}$ is an outcome variable. The target estimand is the outcome mean

$$\psi^{\scriptscriptstyle{\mathsf{MEAN}}}(Q_{O,X}) := \int O \mathrm{d} Q_O$$

Under the sensitivity model (2), the target functional with sensitivity parameter γ is

$$\psi^{\text{MEAN}}(Q_{O,X}^{\gamma}) = \int \frac{\int O \exp(\gamma O) dP_{O|X}}{\int \exp(\gamma O) dP_{O|X}} Q_X,$$
(5)

The SLOPE can be calculated via calculus:

$$\mathsf{SLOPE}(Q_{O,X}^0,\psi^{\mathsf{MEAN}}) = \frac{\partial \psi^{\mathsf{MEAN}}(Q_{O,X}^\gamma)}{\partial \gamma} \bigg|_{\gamma=0} = \mathsf{E}_{Q_X}\{\mathsf{var}_{P_{O|X}}(O \mid X)\} := \mathsf{E}_{Q_X}\{\sigma^2(X)\}.$$

For a specific target (i.e., Q_X), the choice of the $P_{O|X}$ determines the magnitude of $\sigma^2(X)$, and therefore determines the magnitude of SLOPE.



INTUITION: DIRECTION OF SENSITIVITY

$$SLOPE(Q_{O,X}^{0},\psi) = \iint \underbrace{\mathsf{IF}(O,X,\psi(Q_{O,X}^{0}))}_{\text{Influence function}} \underbrace{\{O-\mu(X)\}}_{\text{Direction of sensitivity}} dP_{O|X}dQ_X,$$

To understand the "direction of sensitivity", we consider a broader class of tilting,

$$rac{\mathrm{d} Q_{O|X}^\gamma(o,X)}{\mathrm{d} P_{O|X}(o,X)} \propto
ho(o,x,\gamma),$$

where $\rho(o, x, 0) = 1$ and $\int \rho(o, X, \gamma) dP_{O|X}(o, X) \in (0, \infty)$.

INTUITION: DIRECTION OF SENSITIVITY

$$SLOPE(Q_{O,X}^{0},\psi) = \iint \underbrace{\mathsf{IF}(O,X,\psi(Q_{O,X}^{0}))}_{\text{Influence function}} \underbrace{\{O-\mu(X)\}}_{\text{Direction of sensitivity}} dP_{O|X}dQ_X,$$

To understand the "direction of sensitivity", we consider a broader class of tilting,

$$rac{\mathrm{d} Q_{O|X}^\gamma(o,X)}{\mathrm{d} P_{O|X}(o,X)} \propto
ho(o,x,\gamma),$$

where $\rho(o, x, 0) = 1$ and $\int \rho(o, X, \gamma) dP_{O|X}(o, X) \in (0, \infty)$.

Then the SLOPE is

$$\mathsf{SLOPE}(Q_{O,X}^0,\psi) = \iint \mathsf{IF}(D,X,\psi(Q_{O,X}^0)) \underbrace{\left[\dot{\rho}(O,X,0) - \mathsf{E}_{P_{O|X}}\{\dot{\rho}(O,X,0))\}\right]}_{\mathsf{Direction of sensitivity}} dP_{O|X} dQ_X,$$

where $\dot{\rho}$ is the derivative of ρ with respect to γ .

Explanation: why does Honduras have the lowest SLOPE?

- Mathematically, $\sigma^2(X)$ is lower in Honduras.
 - More heterogeneity in the source population ⇒ more sensitive to violations of the transportability assumption.
 - It matches existing understanding about ease of generalizability when the treatment effect is homogeneous (Tipton and Olsen 2018).



Context-wise, treated households can choose an asset at their choice. In Honduras treated housholds made less diverse choice of the asset.

Table. Most common choice of asset transfer among countries.

Ethiopia	Ghana	Honduras	India	Pakistan	Peru
Sheep & goats (62%)	Goats & hens (44%)	Chickens (83%)	Goats (52%)	Goats (56%)	Guinea pigs (64%)

For all three outcome variables and all (source, target) pairs, we find the median to be in general less sensitive than the mean, i.e., $SLOPE(Q^0_{Y(1),X}, \psi^{MED}) < SLOPE(Q^0_{Y(1),X}, \psi^{MEAN})$.

To investigate what if the data generation changes slightly, we consider a semi-synthetic simulation.

A simplified scenario with binary $X \in \{1, 2\}$ and Gaussian $P_{Y(1)|X}$:

Source:
$$P_{Y(1)|X=1} \sim N(0, 0.5^2)$$
, $P_{Y(1)|X=2} \sim N(\mu_2, \sigma_2^2)$,
Target: $q_1 = Q_X(X=1)$, $q_2 = Q_X(X=2)$

where μ_2 , σ_2^2 , q_1 , q_2 are estimated from data.

- Source $P_{Y(1)|X}$: Peru.
- Target (Q_X): India ($q_1 = 0.2, q_2 = 0.8$) and a synthetic country ($q_1 = q_2 = 0.5$).
- Outcome: per capita consumption.

Peru (P) \Rightarrow India (Q), with $P_{Y(1)|X=1} \sim N(0, 0.5^2), P_{Y(1)|X=2} \sim N(\mu_2, \sigma_2^2).$

Peru (*P*) \Rightarrow India (*Q*), with $P_{Y(1)|X=1} \sim N(0, 0.5^2)$, $P_{Y(1)|X=2} \sim N(\mu_2, \sigma_2^2)$. Question: how would the choice of estimand change if the data changes a bit?

Peru (*P*) \Rightarrow India (*Q*), with $P_{Y(1)|X=1} \sim N(0, 0.5^2), P_{Y(1)|X=2} \sim N(\mu_2, \sigma_2^2)$.

Question: how would the choice of estimand change if the data changes a bit?

- We vary the variance σ_2^2 and calculate the SLOPE in each case.
- For both mean and median, the SLOPE is increasing with σ_2^2 .
- SLOPE for median \leq SLOPE for the mean, where "=" happens when $\sigma_2 = \sigma_1$. Answer: we still choose the median.



Change variance in P_{YIX}

Peru (*P*) \Rightarrow India (*Q*), with $P_{Y(1)|X=1} \sim N(0, 0.5^2)$, $P_{Y(1)|X=2} \sim N(\mu_2), \sigma_2^2)$. Question: how would the choice of estimand change if the data changes a bit?

• We vary the mean μ_2 and calculate the SLOPE in each case.

Peru (*P*) \Rightarrow India (*Q*), with $P_{Y(1)|X=1} \sim N(0, 0.5^2)$, $P_{Y(1)|X=2} \sim N(\mu_2), \sigma_2^2$). Question: how would the choice of estimand change if the data changes a bit?

- We vary the mean μ_2 and calculate the SLOPE in each case.
- When $q_1 = q_2 = 0.5$, SLOPEs do not depend on μ_2 :

$$\mathsf{SLOPE}(Q_{Y(1),X}^{\gamma},\psi^{\mathsf{MEAN}}) = (\sigma_1^2 + \sigma_2^2)/2 \ge \sigma_1 \sigma_2 = \mathsf{SLOPE}(Q_{Y(1),X}^{\gamma},\psi^{\mathsf{MED}}).$$



Change mean in P_{YIX}

Peru (*P*) \Rightarrow India (*Q*), with $P_{Y(1)|X=1} \sim N(0, 0.5^2)$, $P_{Y(1)|X=2} \sim N(\mu_2), \sigma_2^2$). Question: how would the choice of estimand change if the data changes a bit?

- We vary the mean μ_2 and calculate the SLOPE in each case.
- When $q_1 = 0.2$, recall that SLOPEs for mean and median are both weighted average of σ_1^2 and σ_2^2 :

SLOPE $(Q_{Y(1),X}^{\gamma}, \psi^{\text{MEAN}}) = q_1 \sigma_1^2 + q_2 \sigma_2^2$ does not change with μ_2 SLOPE $(Q_{Y(1),X}^{\gamma}, \psi^{\text{MED}}) = w_1 \sigma_1^2 + w_2 \sigma_2^2 = \sigma_2^2 + w_2 \underbrace{(\sigma_2^2 - \sigma_1^2)}_{\geq 0}$, increases with μ_2 .



10 / 10